

Total number of printed pages-7
3 (Sem-6/CBCS) MAT HC 2

2022

MATHEMATICS

(Honours)

Paper : MAT-HC-6026

(Partial Differential Equations)

Full Marks : 60

Time : Three hours

**The figures in the margin indicate
full marks for the questions.**

1. Answer **any seven** : $1 \times 7 = 7$

(i) The equation of the form

$P_p + Q_q = \mathbb{R}$ is known as

(a) Charpit's equation

(b) Lagrange's equation

(c) Bernoulli's equation

(d) Clairaut's equation

(Choose the correct answer)

(ii) How many minimum no. of independent variables does a partial differential equation require?

(iii) Find the degree and order of the equation

$$\frac{\partial^3 z}{\partial x^3} + \left(\frac{\partial^3 z}{\partial x \partial y^2} \right)^2 + \frac{\partial z}{\partial y} = \sin(x + 2y)$$

(iv) Which method can be used for finding the complete solution of a non-linear partial differential equation of first order

(a) Jacobi method

(b) Charpit's method

(c) Both (a) and (b)

(d) None of the above

(Choose the correct answer)

(v) State True Or False :

The equation

$$u_{xx} + u_{yy} + u_{zz} = 0$$

is an Hyperbolic equation.

(vi) Fill in the blanks :

$$\left(\frac{\partial z}{\partial x} \right)^2 + 2 \frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} + z = 0$$

is a _____ order partial differential equation.

(vii) The characteristic equation of $y u_x + x u_y = u$ is

(a) $\frac{dx}{x} = \frac{dy}{y} = \frac{du}{u}$

(b) $\frac{dx}{y} = \frac{dy}{x} = \frac{du}{u}$

(c) $\frac{dx}{u} = \frac{dy}{x} = \frac{du}{y}$

(d) None of the above

(Choose the correct answer)

(viii) State True Or False

$x u_x + y u_y = u^2 + x^2$ is a semi-linear partial differential equation.

(ix) Fill in the blanks :

A solution $z = z(x, y)$ when interpreted as a surface in 3-dimensional space is called _____

(x) The partial differential equation is elliptical if

(a) $B^2 - 4AC > 0$

(b) $B^2 - 4AC \geq 0$

(c) $B^2 - 4AC \leq 0$

(d) $B^2 - 4AC < 0$

(Choose the correct answer)

2. Answer **any four** : $2 \times 4 = 8$

(i) Define quasi-linear partial differential equation and give *one* example.

(ii) Show that a family of spheres

$(x-a)^2 + (y-b)^2 = r^2$ satisfies the partial differential equation

$$z^2(p^2 + q^2 + 1) = r^2$$

(iii) Eliminate the constants a and b from

$$z = (x+a)(y+b).$$

(iv) Determine whether the given equation is hyperbolic, parabolic or elliptic

$$u_{xx} - 2u_{yy} = 0.$$

(v) Solve the differential equation $p + q = 1$.

(vi) Explain the essential features of the "Method of separation of variables".

(vii) Mention when Charpit's method is used. Name a disadvantage of Charpit's method.

(viii) What is the classification of the equation

$$u_{xx} - 4u_{xy} + 4u_{yy} = e^y$$

3. Solve **any three** : $5 \times 3 = 15$

(i) Form a partial differential equation by eliminating arbitrary functions f and F from $y = f(x-at) + F(x+at)$.

(ii) Solve

$$y^2 p - xyq = x(z - 2y)$$

(iii) Find the integral surface of the linear partial differential equation

$$x(y^2 + z)p - y(x^2 + z)q = (x^2 - y^2)z$$

which contains the straight line

$$x + y = 0, z = 1.$$

(iv) Find the solution of the equation $z = pq$ which passes through the parabola

$$x = 0, y^2 = z.$$

(v) Find a complete integral of the equation

$$x^2 p^2 + y^2 q^2 = 1.$$

(vi) Reduce the equation $yu_x + u_y = x$ to canonical form and obtain the general solution.

(vii) Apply the method of separation of variables $u(x, y) = f(x)g(y)$ to solve the equation $u_x + u = u_y$,

$$u(x, 0) = 4e^{-3x}.$$

(viii) Determine the general solution of

$$4u_{xx} + 5u_{xy} + u_{yy} + u_x + u_y = 2.$$

4. Answer **any three**: $10 \times 3 = 30$

(i) Solve $(p^2 + q^2)y - qz = 0$ by Jacobi method.

(ii) Solve $z^2 = pqxy$ by Charpit's method.

(iii) Find the general solution of the differential equation

$$x^2 \frac{\partial z}{\partial x} + y^2 \frac{\partial z}{\partial y} = (x + y)z$$

(iv) Solve $(mz - ny)p + (nx - lz)q = ly - mx$

(v) Use $v = \ln u$ and $v = f(x) + g(y)$ to solve the equation

$$x^2 u_x^2 + y^2 u_y^2 = u^2.$$

(vi) Find the solution of the equation

$$z = \frac{1}{2}(p^2 + q^2) + (p - x)(q - y)$$

which passes through the x axis.

(vii) Find the canonical form of the equation

$$y^2 u_{xx} - x^2 u_{yy} = 0.$$

(viii) Classify the second order linear partial differential equation with example.